

The Nappi-Witten example and gravitational waves

C. DUVAL ⁽¹⁾ Z. HORVÁTH ⁽²⁾ P. A. HORVÁTHY ⁽³⁾

Abstract

The vanishing of the anomaly in the recent example of Nappi and Witten, constructed from the Wess-Zumino-Witten model based on a certain non-semisimple group, follows from a more general result valid for gravitational waves. The construction of the metric is explained.

March 1994.

Final Version (0.111)

submitted to the *Phys. Rev. Lett.* as a *Comment*

⁽¹⁾ Département de Physique, Université d'Aix-Marseille II and Centre de Physique Théorique CNRS-Luminy, Case 907, F-13288 MARSEILLE, Cedex 09 (France)
e-mail:duval@marcptsu1.univ-mrs.fr.

⁽²⁾ Institute for Theoretical Physics, Eötvös University, H-1088 BUDAPEST, Puskin u. 5-7 (Hungary) e-mail: zalanh@ludens.elte.hu

⁽³⁾ Département de Mathématiques, Université de Tours, Parc de Grandmont, F-37200 TOURS (France) e-mail: horvathy@univ-tours.fr

Recently, Nappi and Witten [1] presented a new example of anomaly-free string propagation on four-dimensional space-time with Lorentz signature. Their example, constructed from an ungauged WZW (Wess-Zumino-Witten) model based on a certain non-semisimple group of the type considered before by Cangemi and Jackiw [2], has space-time metric and antisymmetric field

$$d\vec{x}^2 - 2du\left[dv + \frac{1}{2}\epsilon_{ij}x^j dx^i\right] + b du^2 \quad \text{and} \quad b_{ij} = u \epsilon_{ij}, \quad (1)$$

($i = 1, 2$) respectively. Now the metric (1) is a particular case of Brinkmann's gravitational pp wave with a covariantly constant null vector [3],

$$g_{ij}(\vec{x}, u)dx^i dx^j - 2du\left[dv + A_i(\vec{x}, u)dx^i\right] + K(\vec{x}, u)du^2. \quad (2)$$

String propagation in a gravitational wave background has been widely studied [4]. In Ref. [5] — motivated precisely by the WZW model — it is shown, in particular, that for $D = 26$ and for the choice $g_{ij} = g_{ij}(u)$, $A_i = \frac{1}{2}A_{ij}(u)x^j$, $b_{iu} = \frac{1}{2}B_{ij}(u)x^j$, the anomaly vanishes at all loop orders provided the 1-loop condition

$$\triangle K = \frac{1}{2}(A_{ij}A^{ij} - B_{ij}B^{ij}) \quad (3)$$

is satisfied. Then the vanishing of the anomaly in the Nappi-Witten case readily follows since the metric (1) corresponds to $g_{ij} = \delta_{ij}$, $A_{ij} = \epsilon_{ij}$ and $K = b$, and the Nappi-Witten antisymmetric field is brought by the gauge transformation $b_{\mu\nu} \rightarrow b_{\mu\nu} + 2\partial_{[\mu}\lambda_{\nu]}$ with $\lambda_i = \frac{1}{2}u \epsilon_{ij}x^j$ to the form (2) with $B_{ij} = -\epsilon_{ij} = -A_{ij}$. Eq. (3) is identically satisfied.

The result can also be derived from those in Ref. [4]. A rotation with angle $u/2$ in the plane carries in fact the metric (1) into

$$d\vec{x}^2 - 2dudv + K du^2, \quad K = b - \vec{x}^2/4, \quad (4)$$

for which the vanishing of the anomaly at all loop orders has been shown in Ref. [4] under the sole condition $\triangle K + \frac{1}{2}B_{ij}B^{ij} = 0$, which is clearly the same as our (3). Absorbing the constant b into the 'vertical' coordinate as $v \rightarrow v - b(u/2)$ yields furthermore an exact plane wave, $d\vec{x}^2 - 2dudv - (1/4)\vec{x}^2 du^2$, for which the vanishing of anomaly has been proved also non-perturbatively [4].

The symmetries of an exact plane wave are easy to describe. For example, the three commuting symmetries noticed by Nappi and Witten can be made manifest by applying the coordinate transformation $u = \tilde{u}$, $\vec{x} = \sin(u/2)\vec{\tilde{x}}$, $v = \tilde{v} + (1/8)\sin u \vec{\tilde{x}}^2$, which brings the exact plane wave metric into the form $\sin^2(\tilde{u}/2)d\vec{\tilde{x}}^2 - 2d\tilde{u}d\tilde{v}$, for which the three translations $\vec{\tilde{x}} \rightarrow \vec{\tilde{x}} + \vec{\xi}$, $\tilde{v} \rightarrow \tilde{v} + \nu$ are obvious isometries [6]. Let us mention that the exact plane wave — and hence the Nappi-Witten metric (1) — is conformal to Minkowski space $d\vec{X}^2 - 2dUdV$ with conformal factor $\Omega^2(u) = \cos^{-2}(u/2)$, namely through $\vec{X} = \vec{x}/\cos(u/2)$, $U = 2\tan(u/2)$, $V = v + (1/4)\vec{x}^2 \tan(u/2)$.

We now explain the arising of the plane-wave metric on the group. The group of isometries of the Euclidian plane, $\text{ISO}(2) = \text{SO}(2) \ltimes \mathbf{R}^2$, is also a group of canonical transformations for the symplectic structure $dx \wedge dy$. But the $\mathbf{R}$ -bundle $H(1)$ over the plane, endowed with the 1-form $\omega = \frac{1}{2}(xdy - ydx) + dv$ is identified with the Heisenberg group. The ω -preserving transformations of $H(1)$ that cover $\text{ISO}(2)$ form the ‘diamond’ — or oscillator — group $G = \text{SO}(2) \ltimes H(1)$, generated by rotations, $J = y\partial_x - x\partial_y - \partial_u$, translations, $P_x = \partial_x - \frac{1}{2}y\partial_v$ and $P_y = \partial_y + \frac{1}{2}x\partial_v$, and by the central generator $T = \partial_v$. Thus G centrally extends $\text{ISO}(2)$ and is seen to be precisely the group of Nappi and Witten.

Although non semi-simple, G admits a natural bi-invariant Lorentz metric. Indeed, the components of the left-invariant Maurer-Cartan form of G are $\theta = -du$ (rotation), $\alpha + i\beta = e^{-iu}(dx + idy)$ (translations) and ω (extension) as given above. The combination $\alpha^2 + \beta^2 + 2\theta\omega + b\theta^2$ is a right-invariant metric for any constant b , and it yields the metric of Eq. (1). Moreover, the null vector $T = \partial_v$ is covariantly constant.

Note that the Cangemi-Jackiw group [2] arises in the same manner starting with the Poincaré group $\text{ISO}(1, 1)$.

Acknowledgment We are indebted to Professor R. Jackiw for calling our attention to this problem, and to Professors P. Forgács and L. Palla for discussions.

References

- [1] C. Nappi and E. Witten, Phys. Rev. Lett. **71**, 3751 (1993).
- [2] D. Cangemi and R. Jackiw, Phys. Rev. Lett. **69**, 233 (1992).
- [3] H. W. Brinkmann, Math. Ann. **94**, 119-145 (1925).
- [4] D. Amati and C. Klimčík, Phys. Lett. **B219**, 443 (1989); R. Güven, Phys. Lett. **B191**, 275 (1987); H. de Vega and N. Sanchez, Nucl. Phys. **B317**, 706 (1989); A. A. Tseytlin, Phys. Lett. **288**, 279 (1992); G. T. Horowitz and A. R. Steif, Phys. Rev. Lett. **64**, 260 (1990); Phys. Rev. **D42**, 1950 (1990); A. Steif, Phys. Rev. **D42**, 2150 (1990); G. T. Horowitz, in Proc. VIth Int. Superstring Workshop *Strings'90*, Texas '90, Singapore: World Scientific (1991).
- [5] A. A. Tseytlin, Nucl. Phys. **B390**, 153 (1993); C. Duval, Z. Horváth and P. A. Horváthy, Phys. Lett. **B313**, 10 (1993); Mod. Phys. Lett. **A8**, 3749 (1993).
- [6] G. W. Gibbons, Commun. Math. Phys. **45**, 191 (1975).